

SS2-1 Potelligent Technology as Next Generation High Potency Antibody Technology

○Kenya SHITARA¹

¹Strategic Product Planning Department, Kyowa Hakko Kirin Co.Ltd

More than 20 therapeutic antibodies have been approved in US since the late 1990s, and these agents represent a major new class of drugs. However, there still remains room for improvement in the clinical effect and cost of therapeutic antibodies licensed on the market.

Antibody-dependent cellular cytotoxicity (ADCC) is considered to be a major therapeutic function of antibodies. ADCC requires the presence of oligosaccharides in the Fc region and is sensitive to change in the oligosaccharide structure. We have demonstrated that the removal of fucose from IgG1 oligosaccharides results in a very significant enhancement of ADCC. Many therapeutic antibodies approved or clinical development are produced using CHO cells that express high level of α 1,6-fucosyltransferase and consequently produce highly fucosylated antibodies. Potelligent technology allows the stable production of fucose-free antibodies by a fucosyltransferase-knockout CHO cells. Fucose-free antibodies show dramatically enhanced ADCC *in vitro* and *ex vivo*, and improved *in vivo* activity. Thus, the application of fucose-free antibodies is expected to be a promising approach as next-generation therapeutic antibodies with improved efficacy, even when administered at low doses in humans *in vivo*.